

Provider Led Entity

Appropriateness of Advanced Imaging in Patients with Hip Pain: Bibliography

12/01/2020

American Academy of Orthopaedic Surgeons. Management of hip fractures in the elderly: Evidence-based clinical practice guideline (1st Ed.). Rosemont, IL: American Academy of Orthopaedic Surgeons; 2014.

Annabell L, Master V, Rhodes A, Moreira B, Coetzee C, Tran P. Hip pathology: The diagnostic accuracy of magnetic resonance imaging. *J Orthop Surg Res.* 2018; 13(1):127.

Australian and New Zealand Hip Fracture Registry (ANZHFR) Steering Group. Australian and New Zealand guideline for hip fracture care: Improving outcomes in hip fracture management of adults. Sydney: Australian and New Zealand Hip Fracture Registry Steering Group; 2014.

Beaman FD, von Herrmann PF, Kransdorf MJ, Adler RS, Amini B, Appel M, Arnold E, Bernard SA, Greenspan BS, Lee KS, Tuite MJ, Walker EA, Ward RJ, Wessell DE, Weissman BN. ACR Appropriateness Criteria[®] Suspected Osteomyelitis, Septic Arthritis, or Soft Tissue Infection (Excluding Spine and Diabetic Foot). *J Am Coll Radiol.* 2017; 14(5S):S326-S337.

Bencardino JT, Stone TJ, Roberts CC, Appel M, Baccei SJ, Cassidy RC, Chang EY, Fox MG, Greenspan BS, Gyftopoulos S, Hochman MG, Jacobson JA, Mintz DN, Mlady GW, Newman JS, Rosenberg ZS, Shah NA, Small KM, Weissman BN. ACR Appropriateness Criteria[®] Stress (Fatigue/Insufficiency) Fracture, Including Sacrum Excluding Other Vertebrae. *J Am Coll Radiol.* 2017; 14(5S):S293-S306.

Bestic JM, Wessell DE, Beaman FD, Cassidy RC, Czuczman GJ, Demertzis JL, Lenchik L, Motamedi K, Pierce JL, Sharma A, Sloan AE, Than K, Walker EA, Yung EY, Kransdorf MJ. ACR Appropriateness Criteria[®] Primary Bone Tumors. *J Am Coll Radiol.* 2020; 17(5S):S226-S238.

Bussieres AE, Taylor JA, Peterson C. Diagnostic imaging practice guidelines for musculoskeletal complaints in adults-an evidence-based approach. Part 1: Lower extremity disorders. *J Manipulative Physiol Ther* 2007; 30(9):684-717.

Chopra A, Grainger AJ, Dube B, Evans R, Hodgson R, Conroy J, Macdonald D, Robinson P. Comparative reliability and diagnostic performance of conventional 3T magnetic resonance imaging and 1.5T magnetic resonance arthrography for the evaluation of internal derangement of the hop. *Eur Radiol.* 2018; 28(3):963-971.

Crim J, Oserowsky A, Layfield LJ, Schmidt RL. Comparison of radiography and histopathologic analysis in the evaluation of hip arthritis. *AJR Am J Roentgenol.* 2019; 213(4):895-902.

Department of Veterans Affairs / Department of Defence. VA/DoD clinical practice guideline for the non-surgical management of hip and knee osteoarthritis (Version 2.0). 2020

Enseki K, Harris-Hayes M, White DM, Cibulka MT, Woehrle J, Fagerson TL, Clohisy JC. Nonarthritic hip joint pain. Clinical practice guidelines linked to the International Classification of Functioning, Disability and Health from the Orthopaedic Section of the American Physical Therapy Association. *J Orthop Sports Phys Ther.* 2014; 44(6):A1-A32.

Frank JM, Harris JD, Erickson BJ, Slikker W, Bush-Joseph CA, Salata MJ, Nho SJ. Prevalence of femoroacetabular impingement imaging findings in asymptomatic volunteers: A systematic review. *Arthroscopy.* 2015; 31(6):1199-1204.

Grace T, Neumann J, Samaan MA, Souza RB, Majumdar S, Link TM, Zhang AL. Using the Scoring Hip Osteoarthritis with Magnetic Resonance Imaging (SHOMRI) system to assess intra-articular pathology in femoroacetabular impingement. *J Orthop Res.* 2018; 36(11):3064-3070.

Griffin DR, Dickenson EJ, O'Donnell J, Agricola R, Awan T, Beck M, Clohisy JC, Dijkstra HP, Falvey E, Gimpel M, Hinman RS, Holmich P, Kassarian A, Martin HD, Martin R, Mather RC, Philippon MJ, Reiman MP, Takla A, Thorborg K, Walker S, Weir A, Bennell KL. The Warwick Agreement on femoroacetabular impingement syndrome (FAI syndrome): An international consensus statement. *Br J Sports Med.* 2016; 50(19):1169-1176.

Haj-Mirzaian A, Eng J, Khorasani R, Raja AS, Levin AS, Smith SE, Johnson PT, Demehri S. Use of advanced imaging for radiographically occult hip fracture in elderly patients: A systematic review and meta-analysis. *Radiology.* 2020; 296(3):521-531.

Hu LB, Huang ZG, Wei HY, Wang W, Ren A, Xu YY. Osteonecrosis of the femoral head: Using CT, MRI and gross specimen to characterize the location, shape and size of the lesion. *Br J Radiol.* 2015; 88(1046):20140508.

Keeney JA, Nunley RM, Adelani M, Mall N. Magnetic resonance imaging of the hip: Poor cost utility for treatment of adult patients with hip pain. *Clin Orthop Relat Res.* 2014; 472(3):787-792.

Mayo-Smith WW, Hara AK, Mahesh M, Sahani DV, Pavlicek W. How I do it: Managing radiation dose in CT. *Radiology.* 2014; 273(3):657-672.

Mintz DN, Roberts CC, Bencardino JT, BAceci SJ, Caird MS, Cassidy RC, Chang EY, Fox MG, Gyftopoulos S, Kransdorf MJ, Metter DF, Morrison WB, Rosenberg ZS, Shah NA, Small KM, Subhas N, Tambar S, Towers JD, Yu JS, Weissman BN. ACR Appropriateness Criteria® Chronic Hip Pain. *J Am Coll Radiol.* 2017; 14(5S):S90-S102.

Murphey MD, Roberts CC, Bencardino JT, Appel M, Arnold E, Chang EY, Dempsey ME, Fox MG, Fries IB, Greenspan BS, Hochman MG, Jacobson JA, Mintz DN, Newman JS, Rosenberg ZS, Rubin DA, Small KM, Weissman BN. ACR Appropriateness Criteria® Osteonecrosis of the Hip. *J Am Coll Radiol.* 2016; 13(2):147-155.

Muheremu A, Ma J, Amudong A, Ma Y, Niyazi M, Ou Y, Ma Y. Positron emission tomography/computed tomography for osseous and soft tissue sarcomas: A systematic review of the literature and meta-analysis. *Mol Clin Oncol*. 2017; 7(3):461-467.

National Institute for Health and Care Excellence (NICE). The management of hip fracture in adults. 2011 [updated 2017]. London: National Clinical Guideline Centre.

Neiman M, Halshtok Neiman O, Aharoni D, Liberman B, Adar E, Eshed I. Magnetic resonance arthrography of the hip: Prevalence of diagnoses not suspected by the referring physician and correlation with clinical examination and pain score. *Acta Radiol*. 2016; 57(5):595-601.

Neumann J, Zhang AL, Schwaiger BJ, Samaan MA, Souza R, Foreman SC, Joseph GB, Grace T, Majumdar S, Link TM. Validation of scoring hip osteoarthritis with MRI (SHOMRI) scores using hip arthroscopy as a standard of reference. *Eur Radiol*. 2019; 29(2):578-587.

Nguyen BJ, Burt A, Baldassarre RL, Smitaman E, Morshedi M, Kao S, Chang EY, Obrzut S. The prognostic and diagnostic value of 18F-FDG-PET/CT for assessment of symptomatic osteoarthritis. *Nucl Med Commun*. 2018; 39(7):699-706.

O'Sullivan G, Carty F, Cronin CG. Imaging of bone metastasis: An update. *World J Radiol*. 2015; 7(8):202-211.

Reiman MP, Thorborg K, Goode AP, Cook CE, Weir A, Holmich P. Diagnostic accuracy of imaging modalities and injection techniques for the diagnosis of femoroacetabular impingement/labral tear: A systematic review with meta-analysis. *Am J Sports Med*. 2017; 45(11):2665-2677.

Ross AB, Lee KS, Chang EY, Amini B, Bussell JK, Gorbachova T, Ha AS, Khurana B, Klitzke A, Mooar PA, Shah NA, Singer AD, Smith SE, Taljanovic MS, Kransdorf MJ. ACR Appropriateness Criteria® Acute Hip Pain – Suspected Fracture. *J Am Coll Radiol*. 2019; 16(5S):S18-S25.

Saied AM, Redant C, Anthonissen J, Audenaert E, Somers J, Bataillie F, Myncke J, Pattyn C. Conventional versus direct magnetic resonance imaging in detecting labral lesions in femoroacetabular impingement – a retrospective multicenter study. *Acta Orthop Belg*. 2019; 85(1):100-106.

Saied AM, Redant C, El-Batouty M, El-Lakkany MR, El-Adl WA, Anthonissen J, Verdonk R, Audenaert EA. Accuracy of magnetic resonance studies in the detection of chondral and labral lesions in femoroacetabular impingement: Systematic review and meta-analysis. *BMC Musculoskelet Disord*. 2017; 18:83.

Sakellariou G, Conaghan PG, Zhang W, Bijlsma JW, Boyesen P, D'Agostino MA, Doherty M, Fodor D, Kloppenburg M, Miese F, Naredo E, Porcheret M, Iagnocco A. EULAR recommendations for the use of imaging in the clinical management of peripheral joint osteoarthritis. *Ann Rheum Dis*. 2017; 76(9):1484-1494.

Scottish Intercollegiate Guidelines Network (SIGN). Management of hip fracture in older people. A national clinical guideline. 2009. Edinburgh: Scottish Intercollegiate Guidelines Network.

Shah KN, Racine J, Jones LC, Aaron RK. Pathophysiology and risk factors for osteonecrosis. *Curr Rev Musculoskelet Med*. 2015; 8(3):201-209.

Wilson MP, Nobbee D, Murad MH, Dhillon S, McInnes MD, Katlariwala P, Low G. Diagnostic accuracy of limited MRI protocols for detecting radiographically occult hip fractures: A systematic review and meta-analysis. *AJR Am J Roentgenol.* 2020; 215(3):559-567.

Yan K, Xi Y, Sasiponganan C, Zerr J, Wells JE, Chhabra A. Does 3DMR provide equivalent information as 3DCT for the pre-operative evaluation of adult hip pain conditions of femoroacetabular impingement and hip dysplasia? *Br J Radiol.* 2018; 91(1092):20180474.

Yang HL, Liu T, Wang XM, Xu Y, Deng SM. Diagnosis of bone metastases: A meta-analysis comparing FDG PET, CT, MRI and bone scintigraphy. *Eur Radiol.* 2011; 21(12):2604-2617.